



King's Research Portal

DOI:

[10.1136/thoraxjnl-2016-208843](https://doi.org/10.1136/thoraxjnl-2016-208843)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Chamberlain Mitchell, S. A. F., Garrod, R., Clark, L., Douiri, A., Parker, S. M., Ellis, J., Fowler, S. J., Ludlow, S., Hull, J. H., Chung, K. F., Lee, K. K., Bellas, H., Pandyan, A., & Birring, S. S. (2017). Physiotherapy, and speech and language therapy intervention for patients with refractory chronic cough: A multicentre randomised control trial. *Thorax*, 72(2), 129-136. <https://doi.org/10.1136/thoraxjnl-2016-208843>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**PHYSIOTHERAPY, SPEECH AND LANGUAGE THERAPY INTERVENTION FOR
PATIENTS WITH REFRACTORY CHRONIC COUGH: A MULTI-CENTRE
RANDOMISED CONTROLLED TRIAL.**

Authors: Chamberlain Mitchell, Sarah Ann Frances^{1 2} PhD, Garrod, Rachel³ PhD , Clark, Lynne⁴ BSc, Douiri, Abdel^{5 6} PhD, Parker, Sean M⁷ MD, Ellis, Jenny⁷, Fowler, Stephen J⁸ MD, Ludlow Siobhan⁹ BMedSci, Hull, James H¹⁰ PhD, Chung, Kian Fan¹⁰ MD, Lee, Kai.K¹ MD, Bellas, H¹¹ MSc Pandyan, Anand² PhD, Birring, Surinder S.¹ MD

Author Affiliations: ¹Division of Asthma, Allergy and Lung Biology, King's College London, London, United Kingdom; ² School of Health and Rehabilitation, Keele University, Keele, United Kingdom.; ³Denmark Hill Campus, King's College London, United Kingdom; ⁴Speech and Language Therapy Department, King's College Hospital, London, United Kingdom. ⁵Department of Primary Care and Public Health Sciences, King's College London, United Kingdom; ⁶ NIHR Biomedical Centre, King's College London, United Kingdom; ⁷ Respiratory Medicine, Northumbria Healthcare NHSFT, North Tyneside General Hospital, North Shields, United Kingdom; ⁸ Centre for Respiratory Medicine and Allergy, Institute of Inflammation and Repair, The University of Manchester and Lancashire Teaching Hospitals NHS Foundation Trust, Preston, United Kingdom; ⁹ Speech and Language Therapy Department, Leighton Hospital, Mid Cheshire Hospitals Trust, Leighton, United Kingdom; ¹⁰ NIHR Respiratory Biomedical Research Unit at the Royal Brompton NHS Foundation Trust and Imperial College London, London, UK. ¹¹ Physiotherapy Department, University College London Hospitals NHS Foundation Trust, London, UK.

CORRESPONDING AUTHOR: Sarah Ann Frances Chamberlain Mitchell; King's College London, Division of Asthma, Allergy and Lung Biology, Denmark Hill, London, SE9 5RS; 01782 734556. Email address s.chamberlain.mitchell@keele.ac.uk.

Keywords: Chronic cough, Physiotherapy, speech and Language Therapy, Quality of life, cough frequency.

Word Count: 3500

What is the key question?

Is Physiotherapy, speech and language therapy intervention (PSALTI) effective for patients with refractory chronic cough?

What is the bottom line?

PSALTI significantly reduced objective cough frequency and significantly improved quality of life when compared to control intervention.

Why read on?

This study is the first multi-centred randomised controlled trial that demonstrates improvements with Physiotherapy and Speech and Language Therapy Intervention (PSALTI) compared to control intervention using objective outcome measures.

Twitter Feed (140 characters including spaces): Physiotherapy, speech and language therapy (PSALTI) reduces cough and improves quality of life in refractory chronic cough patients

AUTHORS' CONTRIBUTIONS

Conception/design of work: RG, SSB, KL, HB; Study Recruitment: SCM, SSB, SP, SF, JH, KC; Assessments/Treatment delivering in the trial: SCM, LC, JE, SL; Data analysis: SCM, AD, AP, SSB; Drafting manuscript: SCM, RG, SSB; Revised manuscript: All.

ABSTRACT

Background Physiotherapy and Speech and Language Therapy are emerging non-pharmacological treatments for refractory chronic cough. We aimed to investigate the efficacy of a Physiotherapy, Speech and Language Therapy Intervention (PSALTI) to improve health related quality of life (HRQoL) and to reduce cough frequency in patients with refractory chronic cough.

Methods In this multi-centre randomised controlled trial, patients with refractory chronic cough were randomised to four weekly 1:1 sessions of either: PSALTI consisting of education, laryngeal hygiene and hydration, cough suppression techniques, breathing exercises and psycho-educational counselling or control intervention consisting of healthy lifestyle advice. We assessed the change in health related quality of life at week 4 with the Leicester Cough Questionnaire (LCQ). Secondary efficacy outcomes included 24-hour objective cough frequency (Leicester Cough Monitor) and cough reflex sensitivity. The primary analysis used an analysis of covariance adjusted for baseline measurements with the intention-to-treat population. This study was registered at UK Clinical Research Network (UKCRN ID 10678)

Findings Between December 2011, and April 2014, we randomly assigned 75 patients who underwent baseline assessment (34 PSALTI and 41 control). In the observed case analysis, HRQoL (LCQ) improved on average by 1.5 (95%CI: 0.21 to 2.85) points more in PSALTI group than with control ($p=0.024$). Cough frequency improved by 41% (95%CI: 36% to 95%) in PSALTI group relative to control ($p=0.030$). The improvements within the PSALTI group were sustained up to three months. There was no significant difference between groups in the concentration of capsaicin causing 5 or more coughs.

Interpretation Greater improvements in HRQoL and cough frequency were observed with PSALTI intervention. Our findings support the use of PSALTI for patients with refractory chronic cough.

100

101

102

INTRODUCTION

Chronic cough, defined as a cough lasting more than eight weeks, [1] is a prevalent disorder in both the community, [2] and secondary care sectors, accounting for up to 20% of respiratory out-patient clinic referrals. [1, 3] The most common causes of cough in a non-smoking patient with a normal chest radiograph and spirometry are asthma, gastro-oesophageal reflux disease, and rhinitis (upper airway cough syndrome). [1, 4, 5] For a significant number of patients, the cough may remain unexplained or refractory to treatment despite extensive investigation and therapeutic trials. [6] Cough is associated with significant physical and psychological morbidity as well as impaired quality of life. [7-9] There are few effective antitussive therapies for refractory chronic cough. [10, 11] Recent studies suggest a potential role for gabapentin, pregabalin, amitriptyline, morphine and P2X3 receptor inhibitors but they are all associated with significant side effects. [12-16]

Non-pharmacological therapies for refractory chronic cough have shown promising results in a few studies and no significant adverse effects. [17] Non-pharmacological therapies are generally delivered by physiotherapists or speech and language therapists and key components include: education, cough suppression techniques including breathing exercises, vocal hygiene and hydration and psycho-educational counselling. [15, 17-23] Vertigan et al, [19] conducted the only randomised controlled trial of a non-pharmacological intervention for refractory chronic cough and found significantly greater improvements in symptoms of cough for speech pathology management compared to control (general healthy lifestyle advice). The benefits of speech pathology management on objectively measured cough frequency, cough reflex sensitivity and health related quality of life (HRQoL) have not been assessed in a controlled clinical trial, limiting the generalisability of the findings. The minimal clinically important difference of the cough symptom score used in this study has not been defined. Furthermore,

the longer term effect of therapy is not known. [17] A recent study by Patel et al [20] investigated cough-suppression physiotherapy for refractory chronic cough in 23 participants and found a significant improvement in cough related quality of life but this study also did not include a control intervention.

This study therefore aimed to assess the effect of an intervention utilising both physiotherapy and speech and language therapy techniques (Physiotherapy, Speech and Language Therapy Intervention, PSALTI) on HRQoL, objective cough frequency, cough reflex sensitivity and cough severity using a randomised controlled design.

METHODS

A multi-centre, single blinded randomised controlled trial was conducted across three hospitals in the UK (King's College Hospital NHS Foundation Trust, Lancashire Teaching Hospitals NHS Foundation Trust and Northumbria Healthcare NHS Foundation Trust). Two further sites, Royal Brompton & Harefield NHS Foundation Trust and Guy's and St Thomas' NHS Foundation Trust) were recruitment only sites and patients were referred to King's College Hospital to receive the intervention. The study was undertaken between December 2011 and April 2014.

Participants and randomisation

Eligible patients were identified as adults with chronic cough (defined as duration greater than 2 months), with normal chest x-ray, minimal sputum production (less than 10ml sputum a day) and who had negative investigations and/or failed treatment trials for asthma, gastroesophageal reflux and, rhinitis as per British Thoracic Society guidelines. [1] Patients were excluded if they: had an upper respiratory tract infection in past four weeks, were taking angiotensin

converting inhibitor (ACE-I) medication, were current smokers, or had a known respiratory disease (such as lung cancer, pneumonia, pulmonary fibrosis, sarcoidosis, pleural effusion, bronchiectasis). Patients were also excluded if they had vocal cord nodules, malignancy, or evidence of active aspiration.

Once participants had given written consent and completed baseline assessments, they were registered into the randomisation service provided by the King's Clinical Trials Unit, King's College London. This prevented foreknowledge of treatment assignment for the study researchers. Group allocation was concealed from participants until they had completed the study and all post-intervention assessments. Patients were block randomised, stratified by age (above and below 50 years old) and gender.

Control intervention

Participants attended weekly sessions and received one to one standardised healthy lifestyle advice from a health care professional (nurse, physiotherapist or speech and language therapist) over four weeks. The control intervention was based on that used in the trial reported by Vertigan et al. [19] The initial session covered general advice on exercise and physical activity, second session dietary and nutritional advice, third session stress management and fourth session relaxation. The material covered in each session was based on healthy lifestyle advised by the United Kingdom Department of Health and National Health Service. [24-26] The sessions were standardised for all sites by using the same written prompts for therapists and educational information. Face to face training was provided for all site therapists who delivered the healthy lifestyle intervention. The duration of all trial sessions was 45 minutes except the initial session which was one hour.

PSALTI intervention

Participants attended weekly sessions, and received one to one treatment from a health care professional (physiotherapist or speech and language therapist) over four weeks. Session durations were the same as for the control group. The intervention was based on previous speech pathology management and cough-suppression physiotherapy studies for refractory chronic cough reported by Vertigan et al and Patel et al respectively (Table 1). [19, 20] The first session focused on educating participants about chronic cough, introduction to laryngeal hygiene and hydration techniques and cough suppression/distraction. The second and third sessions covered cough suppression techniques in more detail including breathing exercises (table 1). Nasal douching or steam inhalations were recommended to patients with nasal congestion. In the third session psycho-educational counselling techniques were covered with the aid of an information booklet developed jointly by the lead researcher and clinical psychologist at the primary research site. The fourth session consisted of reinforcing all aspects of PSALTI. All components of PSALTI were delivered, however the focus and emphasis on individual techniques varied for each participant, determined by the treating therapist. Airway clearance techniques were included in the PSALTI treatment if the participant's sputum production was close to the upper limit of sputum exclusion criteria. The standardisation of treatment between different hospitals was increased by the use of written treatment plans and educational material. All therapists delivering the treatment were trained in PSALTI prior to commencing the study by the main study researcher.

Table 1: PSALTI components

PSALTI component	Technique
Education	Educate patients on the cough reflex, chronic cough and cough reflex hypersensitivity. Explain the negative effects of repeated coughing Educate patients on voluntary control of cough
Laryngeal hygiene and hydration	Increase frequency and volume of water and non-caffeinated drinks Reduce caffeine and alcohol intake Promote nasal breathing
Cough control	Teach patients to identify their cough triggers Teach patients to use cough suppression or distraction techniques at the first sign or sensation of the need or urge to cough. These cough suppression/distraction techniques include: forced swallow, sipping water and sucking sweets. Teach patients breathing exercises: breathing pattern re-education promoting relaxed abdominal breathing pattern technique; pursed lip breathing to use to control cough.
Psycho-educational counselling	Motivate patients, reiterate the techniques and the aims of therapy Behaviour modification: to try to reduce over-awareness of the need to cough Stress and anxiety management

204 Modified from Chamberlain et al [18]

205

206 **Primary Efficacy Endpoint**

207 HRQoL was assessed with the Leicester Cough Questionnaire (LCQ) at week four, the primary
 208 endpoint. [8] The LCQ is a validated 19-item cough-specific health-related quality of life
 209 questionnaire. Overall scores range from three to twenty-one with a higher score indicating a
 210 better HRQoL. The minimal important difference for this outcome is 1·3. [27] Participants
 211 independently completed questionnaires at baseline, at four weeks (after 4th treatment session)

and at three month follow up. Questionnaires were then placed in sealed envelopes to avoid influencing the treating therapist.

Secondary Efficacy Endpoints

Secondary endpoints were assessed at baseline, 4 weeks and 3 months. Objective cough frequency was assessed with the Leicester Cough Monitor (LCM) a validated, objective, automated and ambulatory cough monitoring device. [28] The LCM consists of a MP3 recording device (Phillips 662 MP3 recorder, UK), external microphone and automated cough detection software. The LCM has been used in previous clinical trials of gabapentin and erythromycin. [12, 29] Participants wore the device for 24 hours at baseline, at four weeks (after fourth treatment session) and three month follow up and were instructed to resume their normal daily activities during this time period. The number of coughs per hour (CF_{perhour}) were recorded.

Capsaicin cough challenge was assessed in a subset of the participants (Kings College Hospital Foundation Trust and Northumbria Healthcare NHS Foundation Trust) to measure participants' cough reflex sensitivity at baseline and at four weeks (after fourth treatment session). Doubling concentrations of capsaicin solution ranging from 0 (saline), 0.49 μm to 1000 μm were administered as per European Respiratory Society guidelines. [30] A dose-response capsaicin cough testing method was used. [30] The nebuliser output was set to 0.01mL/breath⁻¹. The test was discontinued when five or more coughs were induced (C5). In addition, the dose that induced two or more coughs (C2) was recorded.

Cough severity in the past 2 weeks was assessed by a visual analogue scale (0-100mm) as per American College of Chest Physicians guidelines. [31] The vocal performance questionnaire

(VPQ), [32] a 12 item tool was used to assess patients' perceived impact on their voice, since a high prevalence of voice disorders in patients with chronic cough has been reported [33]. A score >12 indicates dysphonia. [32] General health and mood was assessed by Short-Form 36 (SF36) and Hospital Anxiety and Depression scale (HADs). [35, 36] HADs is 14-item questionnaire, a score for either subscale \geq eight indicates mild symptoms, \geq 11 moderate and \geq 15 severe. SF-36 generates two summary scores, physical component summary score (PCS) and mental component summary score (MCS); both range from zero to hundred and a higher score indicates better self-reported health. [37]

Ethics and trial registration

All protocols were approved by the London-Chelsea National Research Ethics Service (NRES) Committee (11-LO-0504). All participants provided written informed consent, and the study was registered with the UK Clinical Research Network (UKCRN ID 10678) and ISRCTN (ISRCTN 73039760).

Role of funding source

The funding bodies had no role in study design, collection, analysis and interpretation of data, in the writing of the report or in the decision to submit for publication.

STATISTICAL ANALYSIS AND SAMPLE SIZE

Power calculations for the primary outcome (LCQ score) were performed based on estimates from a previous study, [38] reporting a mean LCQ score in patients with chronic cough of 14.03 (SD: 3.87). [38] Group sample sizes of 33 in each group achieve 80% power with a significance level of 5% to detect a LCQ change of 2.7 (seen in our pilot study). Allowing for a 25% drop out we aimed to recruit 88 patients in total.

262

263 For each of the variables analysed, univariate descriptive statistics were summarised by
264 randomised group to provide an overview of the data. Summary measures for the baseline
265 characteristics of each group were presented as mean and standard deviation for continuous
266 ‘approximate’ normally distributed variables, medians and interquartile ranges for non-
267 normally distributed variables, and frequencies and percentages for categorical variables.
268 Univariate analyses were performed to compare study group using appropriate statistical tests
269 according to the type and the distribution of the data: independent t-test or Mann-Whitney for
270 continuous variable. Cough frequency and capsaicin data were log transformed prior to
271 analysis.

272

273 Primary efficacy analysis, change in LCQ at week 4, was based on analysis of covariance
274 (ANCOVA) adjusted for the baseline LCQ measurements. The ANCOVA analysis was
275 repeated to adjust for centre and speciality of treating therapist. The analysis used data from the
276 intention-to-treat basis (ITT) population, which included all randomised participants who had
277 received at least one treatment session. In this analysis only observed data were included and
278 no imputation was used for missing data. We also performed an analysis on a per-protocol
279 population (PP) which included patients who completed end of treatment (week 4) cough
280 assessments and who did not deviate from the protocol (established before unmasking).
281 Sensitivity analyses were performed for missing data according to different predefined
282 populations using ANCOVA, with multiple imputations (Online appendix - Table 1). [39]
283 Similar sensitivity analyses were also performed for objective log-transformed cough
284 frequency endpoints (Online appendix - Table 2).

285

The secondary efficacy analysis used data from the intention-to-treat population. In these analyses ANCOVA was used adjusting for baseline variables and only observed data were included without imputation for missing data. A p-value of less than 0.05 was considered statistically significant. All analyses were made using STATA version 12 software (StataCorp LP, College Station, TX).

RESULTS

Participants

Seventy-five participants were randomised and had baseline assessments. One additional patient was randomised to the PSALTI group but did not attend baseline assessments. Four participants did not receive any treatment (PSALTI group (n=3): myocardial infarction prior to treatment, unable to travel to hospital and insufficient time for the study; control group (n=1): undisclosed illness prior to start of treatment). The intention to treat population for LCQ primary analysis consisted of 71 participants (Figure 1 and Online Appendix - Table 1). A total of four participants in the control group and eight participants in the PSALTI group did not receive or complete all treatments for reasons stated in Figure 1. Forty-nine participants completed three month follow up. The consort study flow is described in Figure 1. The baseline characteristics of the randomised participants are described in Table 2. The groups were well-matched with the exception of SF36 Physical Component Summary Scores (higher in the control group).

Table 2. Baseline demographic and clinical characteristics of randomised study participants.

Characteristic	Control (n=41)	PSALTI (n=34)	p value
Age (years)	56 (48 to 67)	61 (53 to 67)	0.239
Female, n (%)	26 (63)	25 (71)	0.459
Cough duration (months)	48 (24 to 126)	60 (30 to 126)	0.279
FEV1 (L, observed), mean(SD)	2.7 (0.9)	2.6 (0.7)	0.517
FEV1/FVC (%), mean(SD)	76 (8.2)	76 (5.0)	0.686
LCQ, mean(SD)	11.9 (3.5)	10.4 (3.6)	0.073
Cough Severity VAS	65 (40 to 83)	63 (49 to 75)	0.652
SF-36 PCS,	47.1 (41.7 to 53.6)	41.1 (35.6 to 49.1)	0.033*
SF-36 MCS	47.7 (38.3 to 54.9)	49.9 (40.5 to 57.0)	0.763
HADs –Anxiety	7 (3 to 10)	7 (4 to 10)	0.785
HADs – Depression	4 (1 to 8)	5 (2 to 6)	0.620
VPQ	17 (11 to 22)	21 (13 to 27)	0.158
CF _{perhour} #	17.0 (0.4)	17.0 (0.4)	0.983
C2 (µm) #	4.01(0.69)	4.74 0.62)	0.677
C5 (µm) #	9.33 (0.56)	8.25 (0.51)	0.708

Data presented as median (IQR) unless otherwise stated.

*p<0.05, # Geometric mean (log SD)

FEV₁, forced expiratory volume in 1 second; FVC, forced vital capacity; LCQ, Leicester cough questionnaire; VAS, visual analogue scale; SF-36, short-form 36 questionnaire; PCS, Physical component score; MCS, Mental component score; HADs, hospital anxiety and depression scale; VPQ, vocal performance questionnaire; CF_{perhour}, cough frequency per hour over a 24 hour period; C2, capsaicin cough challenge – concentration that resulted in two or more coughs; C5, capsaicin cough challenge – concentration that resulted in five or more coughs.

HRQoL - Primary Efficacy Endpoint

There was an improvement in the mean total LCQ score at four weeks with PSALTI; baseline 10.4 vs. four-weeks 14.4, mean difference 3.4, p<0.001. This improvement was larger than that in the control group; baseline mean 11.9 vs. 13.4 at four-weeks, mean difference 1.66, p<0.001 (Online Appendix – Table 3). Total LCQ score at four weeks improved by a mean 1.53 (95% CI 0.21 to 2.85) units more in the PSALTI group than in control (p=0.024), table 3. When adjusted for centre and speciality of therapist, the LCQ score at four weeks improved by a mean

326 of 1.53 (95% CI 0.20 to 2.86), $p=0.024$. The improvement in LCQ with PSALTI was consistent
327 in the per-protocol and sensitivity analyses (Online Appendix - Table 1). The LCQ
328 improvement was sustained from week 4 to 3 months for both groups but there was no
329 significant difference between groups at 3 months (table 3). The LCQ scores and within group
330 differences are presented in online supplement Table 3 and Table 4 respectively. Primary
331 outcome LCQ data (baseline or week 4) was missing in 6.7% of participants. There were no
332 adverse or serious adverse events reported for both interventions.

Table 3. Primary and Secondary efficacy endpoint analysis: Change between PSALTI and control groups at baseline to four weeks and four weeks to three month follow up

	Between group difference		Between group difference	
	Baseline to four weeks		Four weeks to three month follow up	
	Mean Difference (95% CI)	p value	Mean Difference (95% CI)	p value
LCQ Total	1.53 (0.21 to 2.85)	0.024	0.01 (-1.62 to 1.64)	0.994
CF_{perhour} (fold change)	0.59 (0.36 to 0.94)	0.030	1.01 (0.55 to 1.86)	0.966
VAS severity	-9.72 (-20.80 to 1.36)	0.084	1.6 (-15.48 to 18.74)	0.848
SF36 PCS	0.56 (-2.52 to 3.64)	0.717	0.48 (-3.27 to 3.37)	0.977
SF36 MCS	0.81 (-3.10 to 4.72)	0.680	0.72 (-3.06 to 4.51)	0.703
VPQ	3.90 (-0.33 to 8.12)	0.070	-0.20 (-3.43 to 3.03)	0.901
HADS – Anxiety	-0.42 (-1.96 to 1.13)	0.590	0.88 (-0.57 to 2.34)	0.225
HADS – Depression	-0.44 (-1.69 to 0.81)	0.486	-0.18 (-1.36 to 0.99)	0.753
C2 (fold change)	1.11 (0.76 to 1.61)	0.575	NA	NA
C5 (fold change)	1.11 (0.80 to 1.54)	0.512	NA	NA

Between group differences were calculated using ANCOVA adjusted for baseline values.

Positive change in LCQ, SF36 PCS, SF36 MCS, and C5 indicates improvement in symptoms. Negative change in VAS, VPQ, HADS indicates improvement in symptom.

LCQ, Leicester cough questionnaire; CF_{perhour}, cough frequency per hour over a 24 hour period; VAS, visual analogue scale; SF-36, short-form 36 questionnaire; PCS, Physical component score; MCS, Mental component score; VPQ, vocal performance questionnaire HADS, hospital anxiety and depression scale; C2, concentration of capsaicin that caused ≥ 2 coughs, C5, concentration of capsaicin that caused ≥ 5 coughs, NA – not assessed at this time point.

*p<0.05

Objective Cough Frequency

There was a greater reduction in cough frequency after four weeks of treatment in the PSALTI group; geometric mean (SD) 17·0 (2·4) to 9·0 (3·3) coughs per hour ($p=0·002$) versus 17·0 (2·3) to 16·0 (2·2) coughs per hour after control ($p=0·205$), table 4. The control-adjusted improvement in cough frequency per hour in PSALTI was 41% (95% CI [36-95%], $p=0·030$, ANCOVA) at 4 weeks in the primary ITT analysis, (table 3). This improvement was also sustained at three months (Figure 2). The reduction in cough frequency with PSALTI was consistent in per-protocol and sensitivity analyses (Online Appendix - Table 2).

Other questionnaire data

There were no significant between group differences for change (week four minus baseline) in VPQ, depression, anxiety or SF36 (Table 3). There was a greater reduction in VAS cough severity in the PSALTI group compared to control ($p=0·084$, Table 3). There was a reduction in cough severity VAS within groups between week 4 to baseline; control $p=0·007$, PSALTI $p<0·001$ (Table 4).

Table 4: Primary and Second Efficacy Endpoints : within group change

	Change from baseline to four weeks				Change from four weeks to three month follow up			
	PSALTI	P	Control	P	PSALTI	P	Control	P
	Mean Difference (95% CI)		Mean Difference (95% CI)		Mean Difference (95% CI)		Mean Difference (95% CI)	
LCQ Total mean	3.40 (2.26 to 4.55)	0.001*	1.66 (0.78 to 2.54)	<0.001*	-0.17 (-1.49 to 1.15)	0.794	0.27 (-0.82 to 1.35)	0.616
CF_{perhour} (fold change)	0.55 (0.33 to 0.75)	0.002	0.82 (0.60 to 1.22)	0.2053	1.26 (0.84 to 1.90)	0.236	0.91 (0.59 to 1.39)	0.655
VAS severity	-21.18 (-29.83 to -12.53)	< 0.001*	-11.84 (-20.11 to -3.57)	0.007*	9.74 (-3.60 to 23.08)	0.143	0.79 (-10.73 to 12.31)	0.888
C2 (fold change)	1.28 (0.96 to 1.71)	0.089	1.06 (0.81 to 1.36)	0.666	NA	NA	NA	NA
C5 (fold change)	1.24 (1.02 to 1.50)	0.035*	1.08 (0.87 to 1.36)	0.469	NA	NA	NA	NA
SF36 PCS	1.62 (-0.96 to 4.21)	0.208	0.50 (-1.30 to 2.31)	0.574	0.54 (-1.82 to 2.89)	0.639	0.76 (-1.66 to 3.18)	0.522
SF36 MCS	0.53 (-2.69 to 3.75)	0.736	-0.26 (-2.92 to 2.40)	0.843	1.09 (-1.91 to 4.09)	0.456	0.49 (-2.35 to 3.32)	0.727
VPQ	4.04 (0.12 to 7.97)	0.044*	0.73 (-1.94 to 3.39)	0.582	-1.63 (-4.17 to 0.91)	0.193	-0.57 (-3.29 to 2.15)	0.666
HADS – Anxiety	-1.27 (-2.51 to -0.032)	0.045*	-0.90 (-1.96 to 0.17)	0.095	-0.11 (-1.16 to 0.94)	0.826	0.95 (-0.22 to 2.11)	0.104
HADS – Depression	-0.68 (-1.57 to 0.21)	0.126	-0.21 (-1.11 to 0.69)	0.641	0.06 (-1.42 to 1.53)	0.937	0.05 (-0.66 to 0.76)	0.878

Positive change in LCQ, SF36 PCS, SF36 MCS indicates improvement in symptoms. Negative change in VAS, VPQ and HADS indicates improvement in symptoms. LCQ, Leicester cough questionnaire; CF_{perhour}, cough frequency per hour over a 24 hour period; C2, capsaicin cough challenge concentration that resulted in two or more coughs - C5, capsaicin cough challenge – concentration that resulted in five or more coughs; VAS, visual analogue scale; SF-36, short-form 36 questionnaire; PCS, Physical component score; SF-36 MCS, Mental component score; VPQ, vocal performance questionnaire HADS, hospital anxiety and depression scale. NA – not assessed at this timepoint

Cough reflex sensitivity

Sixty participants (80% of ITT group) underwent capsaicin challenge. No significant differences between groups were observed for C2 ($p=0.575$) or C5 ($p=0.512$) (Table 3). There was a within group reduction in C5 with PSALTI ($p=0.035$) but not with control ($p=0.469$), table 4.

DISCUSSION

This study evaluated the efficacy of a physiotherapy and speech and language therapy intervention for patients with refractory chronic cough in a randomised controlled trial. There was a clinically and statistically significant improvement in health-related quality of life with PSALTI compared to control intervention. This was supported by a significant reduction in cough frequency measured objectively. The improvement in health-related quality of life was sustained at a three-month follow-up visit. There was no significant change in cough reflex sensitivity between groups.

Our findings represent an advance from those reported in an earlier study by Vertigan et al. [19] Our study is the first multi-centre trial reported in chronic cough and has the potential to provide the evidence base for access to therapy. Vertigan et al reported a statistically significant reduction in cough symptoms scores but did not include HRQoL or objective assessment with cough frequency monitors. [19] In contrast we assessed HRQoL, objective cough frequency, cough severity VAS and cough reflex sensitivity. We were also able to demonstrate both a clinically and statistically significant improvement in our primary endpoint because the minimally important difference (MID) of the LCQ has been defined. [27] We have shown that

the benefits of PSALTI are sustained after discontinuation of therapy, in contrast to Vertigan et al who did not report follow up data for their participants. [19] One of the strengths of our study was the involvement of multiple centres, the use of standardised treatment protocols and the inclusion of both physiotherapists and speech and language therapists delivering the treatment.

HRQoL, as assessed with the Leicester Cough Questionnaire (LCQ) was selected as the primary outcome measure because it is perhaps the most important outcome measure from the patient's perspective. [7, 8, 40] The HRQoL of our patients was severely impaired, affecting physical, psychological and social domains, comparable to that reported in previous studies of refractory chronic cough. [29, 41] The improvement of HRQoL with PSALTI was large, LCQ 3.4 units. This improvement was greater than the MID of the LCQ, 1.3 units, [27] and that reported for Gabapentin therapy in patients with refractory chronic cough (LCQ improvement 2.5 units). [12] The improvement with PSALTI was smaller when adjusted for the change in the control group (LCQ score control group 1.66 units). HRQoL also improved with control intervention, but to a lesser extent than PSALTI. The aim of the control intervention was to provide patients with an equivalent quantity of clinical attention to the PSALTI intervention. This is additional to what most patients with refractory chronic cough would receive as usual care since physiotherapy and speech and language therapy services are not widely available for refractory chronic cough. It is possible that the control intervention had an anti-tussive effect and that the difference between PSALTI and control group may have been larger if compared to usual care (no active treatment). The control intervention was intended to be non-specific but it is possible that some of its components such as stress/anxiety and lifestyle management may have had a positive benefit, particularly on the central sensitisation pathways that regulate cough.

410

411 The improvement in cough frequency assessed objectively with 24-hour cough monitoring
412 supports the improvement in HRQoL with PSALTI occurred because of an actual reduction in
413 coughing. Cough frequency outcome measures are increasingly being used as end-points in
414 clinical trials to validate the efficacy of anti-tussive therapy. [42, 43] The Leicester Cough
415 Monitor (LCM) has been reported to be a valid method of counting coughs objectively. [28,
416 44] An advantage of cough monitors over subjective measures is that they are not susceptible
417 to the patient's or clinician's perception of cough severity. PSALTI intervention was
418 associated with an additional 41% reduction in cough frequency, which can be considered a
419 large change and is comparable to that observed with pharmacotherapy such as the P2X3
420 inhibitor AF-219. [13] The minimal clinically important difference for cough monitor
421 frequency in chronic cough has not been studied. The reduction of cough frequency was
422 comparable to the minimal important difference reported for acute cough. [42] We also
423 assessed cough severity subjectively with VAS. There was a reduction in cough severity with
424 PSALTI compared to control intervention, and the difference approached statistical
425 significance. The reason for the discrepancy in effect size between HRQoL and VAS findings
426 is not clear. A larger study would be needed to confirm whether PSALTI impacts cough
427 assessed with VAS. Despite their widespread use, VAS have been poorly validated in
428 comparison to HRQoL questionnaires and cough monitoring tools, as acknowledged by the
429 American College of Chest Physicians' Cough Guidelines. [31] There were no between group
430 differences in reported voice related problems. We chose the VPQ, a patient reported
431 questionnaire, to assess voice since it has been reported to have excellent internal consistency,
432 repeatability and responsiveness. [34] There are alternative questionnaires available to assess
433 voice such as the voice handicap index and voice symptom scale [45, 46]. A comparison of

these scales by Webb et al concluded all were valid and reliable questionnaires for assessing patient's perceived voice dysfunction [34].

There were no adverse events associated with PSALTI, specifically no episodes of pulmonary infections. Patients with significant sputum production were excluded because of the potential risk of pulmonary infections associated with cough suppression. Longer-term data with PSALTI is required to fully assess its safety. The mechanism by which PSALTI reduces cough is not clear, nor which component of PSALTI is most effective. PSALTI was not associated with a reduction in cough reflex sensitivity assessed with capsaicin when compared to the change in control group. There was however a significant within group reduction in C5 in the PSALTI group which is consistent with studies by Ryan et al and Vertigan et al, who reported a reduction in cough reflex sensitivity with speech pathology management. [21, 15] The Ryan et al and Vertigan et al studies however did not have a control group (no speech pathology management) for comparison. It is possible that we did not find a between group difference in cough reflex sensitivity due to the small sample size of participants that underwent capsaicin cough challenge testing; further studies are needed to investigate this.

We investigated PSALTI in patients with refractory chronic cough. Our patients had a troublesome chronic cough despite numerous investigations and trials of therapy. A refractory chronic cough may also be referred to as idiopathic, difficult to treat, unexplained, sensory neuropathic and vagal neuropathy cough although some differences exist between groups. [47] PSALTI type treatments have only been studied in patients with refractory chronic cough once they have undergone extensive investigations and/or trials of therapy. The role of PSALTI type treatments earlier in the management of such patients has not been explored and needs to be studied. The efficacy of PSALTI is also unknown in other difficult to treat coughs, such as that

associated with lung cancer, idiopathic pulmonary fibrosis and sarcoidosis and this should be investigated. Further studies are needed to explore the optimum frequency and duration of PSALTI and other non-pharmacological treatments.

There were some limitations to our study. The study was single-blinded. It was not possible to blind the treating therapist to the intervention the patient received. The possibility that unconscious bias could have been conveyed to participants during the course of intervention cannot therefore be discounted. Double blinding is not possible in studies of behavioural intervention. The potential bias was minimised by asking patients to complete their primary outcome measures independently from the treating therapist and participants remained blinded until after completion of the final post-intervention outcome measures. Capsaicin cough reflex tests in some patients were performed by the treating therapist but it is unlikely this influenced the outcome since our findings suggest no change with intervention when adjusted for control. Some components of PSALTI were tailored to the individual, according to clinical need. This may be considered both a limitation and a strength since it reduces the uniformity of intervention delivered but reflects real life clinical practice addressing the needs of an individual. Our study did not meet the intended sample size. This may have affected the power of our analyses and undermined the robustness of the results. Despite this, there was a clinically and statistically significant improvement in the primary outcome measure with PSALTI. Thirty-six (22%) of subjects screened were uncontactable or declined to participate. The clinical characteristics of participants recruited were however consistent with previously reported studies of refractory chronic cough [29]. A significant number of patients were lost to follow up for the 3 month visit where secondary endpoints were assessed. This was largely from the control group. There was no significant difference in LCQ between groups at 3 months; this could be a consequence of a smaller sample size or a reduction in the long term

benefit of PSALTI following cessation of therapy. The long term benefits of PSALTI needs to be confirmed in larger studies. It is possible that some of the benefit of PSALTI may have been a consequence of intense supervision. The control intervention was however identical in frequency and duration of visits.

In conclusion, PSALTI is an effective therapy for patients with refractory chronic cough. It is associated with a significant improvement in health-related quality of life, and cough frequency compared to control. The optimal components of PSALTI and number of sessions of therapy need to be determined in future studies. The effectiveness of PSALTI used earlier in the treatment of chronic cough and other patient groups with difficult-to-treat cough, should be evaluated. There is also a need for improved access to physiotherapy and speech and language therapy services for patients with refractory chronic cough.

ACKNOWLEDGEMENTS

This study represents independent research supported by the National Institute for Health Research (NIHR) / Wellcome Trust King's Clinical Research Facility and the NIHR Biomedical Research Centre and Dementia Unit at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. Dr. A. Douiri was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. All authors would like to acknowledge a number of key people who helped with this trial. We would like to thank Dr Nicholas Hart for his help with applying for

NIHR grant funding, Janet Mills and Maureen Armstrong for their help with recruitment of participants and study/data management at Royal Preston and North Tyneside Hospitals respectively, Natasha Muzengi, Rachel Harding, Aliya Kaaba and Katie Pidgeon who all helped with conducting assessments and patient recruitment at King's College Hospital.

Declaration of Interests:

All authors have completed conflict of interest forms. SSB, RG, SL, JE, AD, SCM, SF, SP, KL, HB, JH and LC have all declared no conflict of interests. AP reports grants from NIHR, personal fees from Industry (allergan, Merz and IPSEN), grants and non-financial support from Industry (Allergan and Biometrics Ltd), outside the submitted work. KFC has no disclosure for the work under consideration. Outside this work, he has received honoraria for participating in Advisory Board meetings regarding treatments for asthma and chronic obstructive pulmonary disease for GSK, AZ, Novartis and J&J and has received grant funding through his institution from Pfizer, GSK and Merck.

FUNDING

This work was supported by a grant from the Chartered Society of Physiotherapy Charitable Trust, United Kingdom (Award PRF 10/4). Additional funding was obtained from NIHR-CRN. SB was supported by London National Institute for Health Research (NIHR) / Wellcome Trust King's Clinical Research Facility and the NIHR Biomedical Research Centre and Dementia Unit at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health., UK.

FIGURE LEGENDS

Figure 1. Trial CONSORT flow diagram

Figure 2. Change in objective cough frequency in PSALTI and control groups

REFERENCES

1. Morice, A.H., McGarvey, L., Pavord, I. On behalf of the British Thoracic Society. Recommendations for the management of cough in adults. *Thorax* 2006; **61**: Suppl 1: i1–24.
2. Ford, A.C., Forman, D., Moayyedi, P., Morice, A.H. Cough in the community: a cross sectional survey and the relationship to gastrointestinal symptoms. *Thorax* 2006; **61**:11: 975–9.
3. Morice, A. Chronic cough: epidemiology. *Chron Respir Dis* 2008; **5**:1: 43–7.
4. Morice, A.H., Fontana, G.A., Sovijarvi, A.R., et al., The diagnosis and management of chronic cough. *Eur Respir J* 2004; **24**:3: 481–92.
5. Irwin, R., Baumann, M.H., Bolser, D.C., et al., Diagnosis and management of cough executive summary: ACCP evidence-based clinical practice guidelines. *Chest* 2006; **129**: 1 Suppl: 1S–23S.
6. Haque, R.A., Usmani, O.S., Barnes, P.J. Chronic idiopathic cough: a discrete clinical entity? *Chest* 2005; **127**:5: 1710–3.
7. Brignall, K., Jayaraman, B., Birring, S.S. Quality of life and psychosocial aspects of cough. *Lung* 2008. **186**: Suppl 1: S55–8.

8. Birring, S.S., Prudon, B., Carr, A.J., Singh, S.J., Morgan, M.D., Pavord, I.D.
Development of a symptom specific health status measure for patients with chronic
cough: Leicester Cough Questionnaire (LCQ). *Thorax* 2003. **58**:4: 339–43.
9. French, C.L., Irwin, R.S., Curley, F.J., Krikorian, C.J. Impact of chronic cough on
quality of life. *Arch Intern Med* 1998. **158**:15:1657–61.
10. Chung, K.F. Clinical cough VI: the need for new therapies for cough: disease-specific
and symptom-related antitussives. *Handb Exp Pharmacol* 2009. **187**: 343–68.
11. Belvisi, M.G., Geppetti, P. Cough • 7: Current and future drugs for the treatment of
chronic cough. *Thorax* 2004. **59**:5: 438–440.
12. Ryan, N.M., Birring, S.S., Gibson, P.G. Gabapentin for refractory chronic cough: a
randomised, double-blind, placebo-controlled trial. *Lancet* 2012. **380**:9853:1583–9.
13. Abdulqawi, R., Dockry, R., Holt, K. et al. P2X3 receptor antagonist (AF-219) in
refractory chronic cough: a randomised, double-blind, placebo-controlled phase 2
study. *Lancet*, 2015. **385**:9974: 1198–205
14. Morice, A.H., Monon, M.S., Mulrennan, S.A., et al., Opiate therapy in chronic cough.
Am J Respir Crit Care Med 2007. **175**:4:312–5.

15. Vertigan, A.E., Kapela, S.L., Ryan, N.M., Birring, S.S., McElduff, P., Gibson, P.G. Pregablin and Speech Pathology Combination Therapy for Refractory Chronic Cough: A Randomized Controlled Trial. *Chest* 2016. **149**:3:639-648.
16. Jeyakumar, A., Brickman, T. M., Haben, M. Effectiveness of amitriptyline versus cough suppressants in the treatment of chronic cough resulting from postviral vagal neuropathy. *Laryngoscope* 2006. **116**:12:2108–12.
17. Chamberlain, S., Birring, S.S., Garrod, R. Nonpharmacological interventions for refractory chronic cough patients: systematic review. *Lung* 2014. **192**:1: 75–85.
18. Chamberlain, S., Garrod, R., Birring, S.S. Cough suppression therapy: does it work? *Pulm Pharmacol Ther* 2013. **26**:5:524–7.
19. Vertigan, A.E., Theodoros, D.G., Gibson, P.G., Winkworth, A.L. Efficacy of speech pathology management for chronic cough: a randomised placebo controlled trial of treatment efficacy. *Thorax* 2006. **61**:12:1065–9.
20. Patel, A.S., Watkin, G., Willig, B., et al. Improvement in health status following cough-suppression physiotherapy for patients with chronic cough. *Chron Respir Dis* 2011. **8**:4:253–8.
21. Ryan, N.M., Vertigan, A.E., Bone, S., Gibson, P.G. Cough reflex sensitivity improves with speech language pathology management of refractory chronic cough. *Cough* 2010. **6**:5.

22. Ryan, N.M., Vertigan, A.E., Gibson, P.G. Chronic cough and laryngeal dysfunction improve with specific treatment of cough and paradoxical vocal fold movement. *Cough* 2009. **5**:4.
23. Vertigan, A., Theodoros, D., Winkworth, A., Gibson, P. Chronic cough a tutorial for speech-pathologists. *Journal of medical speech-language pathology. Journal of Medical Speech-Language Pathology* 2007. **15**:3: 189–206.
24. Department of Health (2011) Start Active, Stay Active. UK, Department of Health
25. NHS (2013) Eight tips for healthy eating. Available at <http://www.nhs.uk/Livewell/Goodfood/Pages/eight-tips-healthy-eating.aspx> [Accessed 05/10/2015]
26. NHS (2013) Relaxation tips to relieve stress. Available at <http://www.nhs.uk/Conditions/stress-anxiety-depression/Pages/ways-relieve-stress.aspx> [Accessed 05/10/2015]
27. Raj, A.A., Pavord, D.I., Birring, S.S. Clinical cough IV: what is the minimal important difference for the Leicester Cough Questionnaire? *Handb Exp Pharmacol* 2009. **187**:311–20.
28. Birring, S.S., Fleming, T., Matos, S., Rai, A.A., Evans, D.H., Pavord, I.D. The Leicester Cough Monitor: preliminary validation of an automated cough detection system in chronic cough. *Eur Respir J* 2008. **31**:5:1013–8.

29. Yousaf, N., Monteiro, W., Parker, D., Matos, S., Birring, S., Pavord, I.D. Long-term low-dose erythromycin in patients with unexplained chronic cough: a double-blind placebo controlled trial. *Thorax* 2010. **65**:12:1107–10.
30. Morice, A.H., Fontana, G.A., Belvisi, M.G. et al. ERS guidelines on the assessment of cough. *Eur Respir J* 2007. 29:6:1256–76.
31. Boulet, L., Coeytaux, R.R., McCrory, D.C., et al. Tools for Assessing Outcomes in Studies of Chronic Cough: CHEST Guideline and Expert Panel Report. *Chest* 2015. 147:3:804–14.
32. Carding, P.N., Horsley, I.A. An evaluation study of voice therapy in non-organic dysphonia. *Eur J Disord Commun* 1992. **27**:2:137–58.
33. Vertigan, A. E., Theodoros, D. G., Winkworth, A. L. and Gibson, P. G. Perceptual voice characteristics in chronic cough and paradoxical vocal fold movement. *Folia Phoniatr Logop* 2007. 59:5:256-67.
34. Webb, A. L., Carding, P. N., Deary, I. J., MacKenzie, K., Steen, I. N. and Wilson, J. A. Optimising outcome assessment of voice interventions, I: Reliability and validity of three self-reported scales. *J Laryngol Oto* 2007. 121:8: 763-7.
35. Brazier, J.E., Harper, R., Jones, N.M., et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ* 1992. **305**:6846:160–4.
36. Zigmond, A.S., Snaith, R.P. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983. **67**:6:361–70.

678

679 37. Lacson, E., Xu, J., Lin, S.F., Dean, S.G., Lazarus, J.M., Hakim, R.M. A comparison of
680 SF-36 and SF-12 composite scores and subsequent hospitalization and mortality risks
681 in long-term dialysis patients. *Clin J Am Soc Nephrol* 2010. **5**:2:252–60.

682

683 38. Birring, S.S., Pavord, I.D. Assessment of gender differences in health status with the
684 Leicester Cough Questionnaire (LCQ). *Thorax* 2009. **64**:11:1008–9.

685

686 39. Little, R.J., D’Agostino, R., Cohen, M.L., et al., The prevention and treatment of
687 missing data in clinical trials. *N Engl J Med* 2012. **367**:14:1355–60.

688

689 40. Spinou, A., Birring, S. S. An update on measurement and monitoring of cough: what
690 are the important study endpoints? *J Thorac Dis* 2014. **6**:Suppl 7: S728–34.

691 41. Birring, S.S., Murphy, A.C., Scullion, J.E., Brightling, C.E., Browning, M., Pavord,
692 I.D. Idiopathic chronic cough and organ-specific autoimmune diseases: a case-control
693 study. *Respir Med* 2004. **98**:3:242–6.

694 42. Lee, K.K., Savani, A., Matos, S., Evans, D.H., Pavord, I.D., Birring, S.S. Four-hour
695 cough frequency monitoring in chronic cough. *Chest* 2012.**142**:5:1237–43.

696 43. Lee, K.K., Matos, S., Evans, D.H., White, P., Pavord, I.D., Birring, S.S. A
697 Longitudinal Assessment of Acute Cough. *American Journal of Respiratory and*
698 *Critical Care Medicine* 2013. **187**:9:991–997.

699

700 44. Yousaf, N., Monteiro, W., Matos, S., Birring, S.S., Pavord, I.D. Cough frequency in
701 health and disease. *Eur Respir J* 2013. **41**:1:241–3.

702

703 45. Jacobson, B.H., Johnson, A., Grywalski, C., Silbergleit, A., Jacobson, G. Benninger,
704 M.A., Newman, G.W. The Voice Handicap Index (VHI): Development and
705 Validation. *American Journal of Speech-Language Pathology* 1997. **6**:66-70.

706

707 46. Dreary, I.J., Wilson, J.A., Carding, P.N., MacKenzie, K. VoiSS: a patient-derived
708 Voice Symptom Scale. *J PsychoSom Res* 2003. **54**:5:483-489.


709

710 47. Birring, S.S. Controversies in the evaluation and management of chronic cough. *Am J*
711 *Respir Crit Care Med* 2011. **183**:6:708-15.

712


713

Figure. 1. Trial CONSORT flow diagram



*One additional participant was randomised and withdrew before baseline assessments.

Figure 2: Change in objective cough frequency in PSALTI and control groups



Data presented as Geometric Mean (log 95%CI) coughs per hour. PSALTI: physiotherapy speech and language therapy intervention.

Online Appendix

Supplementary Methods: Sensitivity Analysis

A total of 75 patients were randomised and underwent baseline assessment. One additional patient was randomised but then declined participation in the trial before baseline measures could be assessed. All patients were reviewed on a case-by-case basis prior to data unblinding and analysis, to identify patients for sensitivity analyses.

The following populations were defined for the primary efficacy analysis:

- 1) All randomised patients who had baseline assessment.
- 2) Intention to treat (ITT): all patients who received at least one treatment session.
- 3) All patients who received at least one treatment session and, in addition, had not been withdrawn from study treatment prior to week 4.
- 4) Per protocol: all patients who had both baseline and week 4 measurements and did not deviate from protocol.

Sensitivity analysis was done on predefined populations using analysis of covariance adjusted for baseline measurements ANCOVA, with multiple imputation method for missing data. Multiple imputation method used was based on multivariate normal regression using an iterative Markov chain Monte Carlo (MCMC) method to impute missing values. The imputation process was repeated to create 10 multiple imputed datasets. Missing week 4 Leicester Cough Questionnaire were imputed, as long as baseline data was available. Hence no values were imputed where data was missing for both baseline and week 4 in a period. For cough frequency, data were imputed when week 4 values were missing, in a secondary analysis using ANCOVA, with multiple imputation method.

Online supplement Table 1 and 2 summarise sensitivity analyses results of LCQ and 'log' cough frequency per hour.

Online Supplement Table 1: Sensitivity analyses for LCQ.

Population	Imputation method	N	Estimate	[95% CI]	p-value
Randomised patients*	ANCOVA, with Multiple imputation	75	1.6	0.14 to 2.95	0.032
All patients who received at least one study treatment (ITT)	ANCOVA, with Multiple imputation	71	1.5	0.20 to 2.76	0.024
All patients who received at least one study treatment and had not deviated the protocol	ANCOVA, with Multiple imputation	67	1.4	0.08 to 2.78	0.039
Per Protocol	ANCOVA, no imputation	63	1.5	0.21 to 2.85	0.024

ANCOVA: analysis of covariance analysis, N: sample size; CI: confidence intervals, LCQ: Leicester Cough Questionnaire (cough specific quality of life). ITT: intention to treat

*All randomised patients with baseline assessment (excludes 1 patient randomised who then declined participation and no baseline assessment).

Online supplement Table 2: Sensitivity analyses results of mean fold difference in cough frequency per hour.

Population	Imputation method	N	Estimate	[95% CI]	p-value
Randomised patients*	ANCOVA, with Multiple imputation	69	0.58	0.34 to 0.98	0.043
All patients who received at least one study treatment (ITT)	ANCOVA, with Multiple imputation	66	0.59	0.35 to 1.00	0.049
All patients who received at least one study treatment and had not deviated the protocol	ANCOVA, with Multiple imputation	63	0.57	0.36 to 0.90	0.019
Per Protocol	ANCOVA, no imputation	53	0.59	0.36 to 0.95	0.030

ANCOVA: analysis of covariance analysis, N: sample size; CI: confidence intervals. ITT: intention to treat

*All randomised patients with baseline assessment (excludes one patient who was randomised then decline participation).

Online supplement Table 3: Leicester Cough Questionnaire HRQoL scores.

	PSALTI Group	Control Group
Baseline	10.4 (3.6)	11.9 (3.5)
Four weeks	14.4 (3.3)	13.4 (3.7)
Three Months	14.8 (3.9)	13.6 (3.4)

Data presented as Mean (standard deviation). HRQoL: health related quality of life. PSALTI: physiotherapy and speech and language therapy.